



Reference

Accession

Patient

NHS No

D.O.B.

Patient Ref

Reason

Varicose vein

Outcome

DVT negative, Superficial oedema, Competent

Right			Left	
Deep Veins	Patency	Competency	Patency	Competency
Common Iliac Vein				
External Iliac Vein				
Internal Iliac Vein				
Common Femoral Vein			Widely Patent	Competent
Profunda Vein			Widely Patent	Competent
Superficial Femoral Vein			Widely Patent	Competent
Popliteal Vein			Widely Patent	Competent
Posterior Tibial Vein			Widely Patent	Competent
Anterior Tibial Vein			Widely Patent	Competent
Peroneal Vein			Widely Patent	Competent
Soleal Vein				
Gastrocnemius			Widely Patent	Competent
Superficial Veins				
Saphenofemoral Junction			Widely Patent	Competent
L Saphenous Vein Above			Widely Patent	Competent
L Saphenous Vein Below			Widely Patent	Competent
Vein of Giacomini			Widely Patent	Competent
Saphenopopiteal Junction			Not Identified	
S Saphenous Vein			Widely Patent	Competent
Evidence of D.V.T.				
Above the knee			No	
Popliteal			No	
Below the knee			No	

Notes

LEFT LOWER LIMB VENOUS DUPLEX ASSESSMENT

Iliac veins not viewed. Flow in the common femoral vein is phasic with respiration and a normal response on Valsalva manoeuvre, suggesting proximal vein patency. All visualised deep veins appear widely patent and competent with no evidence of previous DVT.

All truncal, superficial veins appear to be widely patent and competent.