



Reference

Accession

Patient

NHS No

D.O.B.

Patient Ref

Reason

Routine


Outcome

Widely patent


Right

130

1.00



Good




Good



Good

140

1.08



Good

Left

Brachial

Common Femoral

High Thigh

Low Thigh

Popliteal

High Calf

Peroneal

Anterior Tibial

Posterior Tibial

Dorsalis Pedis

Toe Pressure

Post Exercise

Good

Good

Good

140

1.08

Good

Foot Flex

130

1.00

Notes

BILATERAL LOWER LIMB ARTERIAL PRESSURES & WAVEFORMS / ABDOMINAL AORTA ASSESSMENT

Abdominal aorta is widely patent with good bi/triphasic waveforms and PSV 90cm/s. The abdominal aorta appears of normal calibre (maximum AP = 1.6cm), with no evidence of focal dilatation or aneurysm identified.

Good bi/triphasic waveforms detected in the right and left common femoral, popliteal, posterior tibial and

Assessed by

Lukasz Koprowski

Printed on 08/06/2019 at 11:40 am

Checked by



Patient

NHS No

D.O.B.

Patient Ref

anterior tibial arteries.

Bilateral, resting ABPIs are within normal limits, with no significant reduction in systolic ankle pressure observed following a one minute exercise challenge.

CONCLUSION: No evidence of significant arterial disease identified in the right or left lower limbs from this assessment.