






Reason Claudication
Outcome Aneurysm, Occlusion

| Right | Left |
|---|---|
| <div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 2px 10px;">115</div> <div style="border: 1px solid black; padding: 2px 10px;">1.00</div> </div> | <div style="display: flex; justify-content: space-between; align-items: center;"> <div style="width: 60%;"> Brachial Common Femoral High Thigh Low Thigh Popliteal </div> <div style="width: 35%;"> <div style="border: 1px solid black; padding: 2px 10px;">Good</div> <div style="border: 1px solid black; padding: 2px 10px;">Reduced</div> </div> <div style="width: 5%; text-align: center;">   </div> </div> |
| <div style="display: flex; justify-content: space-between; align-items: center;">  <div style="width: 60%;"> <div style="border: 1px solid black; padding: 2px 10px;">Slightly Reduced</div> <div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 2px 10px;">110</div> <div style="border: 1px solid black; padding: 2px 10px;">0.96</div> </div> </div> </div> | <div style="display: flex; justify-content: space-between; align-items: center;"> <div style="width: 60%;"> High Calf Peroneal Anterior Tibial Posterior Tibial </div> <div style="width: 35%;"> <div style="border: 1px solid black; padding: 2px 10px;">Reduced</div> <div style="border: 1px solid black; padding: 2px 10px;">Reduced</div> </div> <div style="width: 5%; text-align: center;">   </div> </div> |
| <div style="display: flex; justify-content: space-between; align-items: center;"> <div style="width: 60%;"> Dorsalis Pedis Toe Pressure </div> <div style="width: 35%;"></div> </div> | <div style="display: flex; justify-content: space-between; align-items: center;"> <div style="width: 60%;"> Post Exercise </div> <div style="width: 35%;"></div> </div> |
| <div style="display: flex; justify-content: space-between; align-items: center;"> <div style="width: 60%;"> <div style="border: 1px solid black; padding: 2px 10px;">Foot Flex</div> <div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 2px 10px;">90</div> <div style="border: 1px solid black; padding: 2px 10px;">0.78</div> </div> </div> </div> | |

Notes

LEFT LOWER LIMB ARTERIAL DUPLEX ASSESSMENT AND AAA SCREEN

The abdominal aorta appears aneurysmal, measuring 4.3cm (outer-outer) and 4.0cm (inner-inner) LS and TS.

CFA - moderate disease with good triphasic waveforms, PSV 221cm/s.

PFA - mild disease with good triphasic waveforms, PSV 143cm/s.

SFA - vessel appears to occlude ~0.7cm from the origin. Flow reforms in the adductor region, with reduced

Assessed by Rachel Johnson

Printed on 13/06/2019 at 11:04 am

Checked by _____

monophasic waveforms, PSV 14cm/s.

POPA - mild disease with reduced monophasic waveforms, PSV 21cm/s. TPT appears patent with evidence of two vessel run-off.

ATA - mild disease with reduced/weak monophasic waveforms, PSV 10cm/s.

PTA - appears patent proximally with mild disease and reduced monophasic waveforms, PSV 16cm/s. No colour flow noted in the mid to distal vessel ?patency.

Resting ABPI not completed due to weak signal on HHD.

CONCLUSION: Evidence of a small AAA. Evidence of left SFA occlusion.

