


Reason Routine
Outcome Widely patent


Right

125

1.00




Good



Good



Good



Good

130

1.04

Brachial

Common Femoral

High Thigh

Low Thigh

Popliteal

High Calf

Peroneal

Anterior Tibial

Posterior Tibial

Dorsalis Pedis

Toe Pressure

Foot Flex

110


0.88

Post Exercise

Foot Flex

100

0.80



Good

120

0.96

Notes

RIGHT LOWER LIMB ARTERIAL DUPLEX ASSESSMENT

CFA - widely patent with good triphasic waveforms, PSV 166cm/s.

PFA - widely patent with good triphasic waveforms, PSV 150cm/s.

SFA - widely patent with good triphasic waveforms, PSV 170-122cm/s.

POPA - widely patent with good triphasic waveforms, PSV 86cm/s. TPT appears patent with evidence of two vessel run-off.

ATA/PTA - appears widely patent with good biphasic and triphasic waveforms, PSV 35cm/s and 79cm/s

Assessed by Rachel Johnson

Printed on 13/06/2019 at 10:56 am

Checked by

respectively.

Bilateral resting ABPIs are within normal limits with no significant reduction post foot-flex exercise.

CONCLUSION: No evidence of any significant right lower limb arterial disease identified from this assessment.