











Reason Ulceration
Outcome Poor images

Right			Left	
	115 1.00	Brachial		
	Good	Common Femoral	Good	
		High Thigh		
		Low Thigh		
	Good	Popliteal	Good	
		High Calf		
		Peroneal	Good	
	Good 140 1.22	Anterior Tibial	Good 140 1.22	
	Good	Posterior Tibial		
	Good	Dorsalis Pedis	Good	
		Toe Pressure		
		Post Exercise		

Notes

BILATERAL LOWER LIMB ARTERIAL DUPLEX ASSESSMENT

*Patient unable to lie flat or lie legs flat - poor access to the groin.

RIGHT

CFA - mild disease with good triphasic waveforms, PSV 115cm/s.

PFA - not visualised.

SFA - mild disease with good triphasic waveforms, PSV 107-124cm/s.

POPA - mild disease with good triphasic waveforms, PSV 106cm/s. TPT appears patent with evidence of

Assessed by Rachel Johnson

Printed on 13/06/2019 at 10:15 am

Checked by

two vessel run-off.

ATA/PTA - only proximal vessel assessed due to calf dressings - good triphasic waveforms, PSV 62cm/s and 76cm/s respectively.

DP - good triphasic waveforms on HHD.

LEFT

CFA - mild disease with good triphasic waveforms, PSV 150cm/s.

PFA - not visualised.

SFA - mild disease with good triphasic waveforms, PSV 116-70cm/s.

POPA - mild disease with good triphasic waveforms, PSV 136cm/s. TPT appears patent with evidence of one vessel run-off.

ATA/PeRoA - only proximal vessel assessed due to calf dressings - good triphasic waveforms, PSV 88cm/s and 60cm/s respectively.

DP - good triphasic waveforms on HHD.

Resting ABPIs are within normal limits - patient unable to complete adequate exercise challenge.

CONCLUSION: Pressures and waveforms suggest no evidence of any significant right or left lower limb arterial disease.

