

Reason Varicose vein
Outcome Competent

| Deep Veins | Right | | Left | |
|---------------------------|----------------|------------|----------------|------------|
| | Patency | Competency | Patency | Competency |
| Common Iliac Vein | | | | |
| External Iliac Vein | | | | |
| Internal Iliac Vein | | | | |
| Common Femoral Vein | Widely Patent | Competent | Widely Patent | Competent |
| Profunda Vein | Widely Patent | Competent | Widely Patent | Competent |
| Superficial Femoral Vein | Widely Patent | Competent | Widely Patent | Competent |
| Popliteal Vein | Widely Patent | Competent | Widely Patent | Competent |
| Posterior Tibial Vein | Patent | | Patent | |
| Anterior Tibial Vein | Patent | | Patent | |
| Peroneal Vein | Patent | | Patent | |
| Soleal Vein | Not Identified | | Not Identified | |
| Gastrocnemius | Patent | | Patent | |
| Superficial Veins | | | | |
| Saphenofemoral Junction | Patent | Competent | Patent | Competent |
| L Saphenous Vein Above | Patent | Competent | Patent | Competent |
| L Saphenous Vein Below | Patent | Competent | Patent | Competent |
| Vein of Giacomini | Patent | Competent | Not Identified | |
| Saphenopopiteal Junction | Not Identified | | Patent | Competent |
| S Saphenous Vein | Patent | Competent | Patent | Competent |
| Evidence of D.V.T. | | | | |
| Above the knee | No | | No | |
| Popliteal | No | | No | |
| Below the knee | No | | No | |

Notes

BILATERAL LOWER LIMB VENOUS DUPLEX ASSESSMENT

Iliac veins not viewed. Flow in the common femoral vein is phasic with respiration and demonstrates a normal response on Valsalva manoeuvre, suggesting proximal vein patency. All visualised deep veins, proximal to and including the popliteal vein appear widely patent and competent with no evidence of previous DVT. Calf veins were difficult to visualise due to depth and patient discomfort during calf augmentation, but appear patent with reasonable colour-filling.

RIGHT

SFJ is competent.

LSV is competent in the thigh and calf.

Assessed by Rachel Johnson

Printed on 18/06/2019 at 11:18 am

Checked by

Transverse (AP) dimensions of LSV: Proximal thigh - 0.64cm, Mid thigh - 0.5cm, Distal thigh - 0.35cm,
Proximal calf - 0.26cm, Mid calf - 0.27cm, Distal calf - 0.28cm

SPJ was not identified.

SSV is competent and is continuous with a competent vein of Giacomini.

LEFT

SFJ is competent.

LSV is competent in the thigh and calf.

Transverse (AP) dimensions of LSV: Proximal thigh - 0.5cm, Mid thigh - 0.43cm, Distal thigh - 0.43cm,
Proximal calf - 0.31cm, Mid calf - 0.29cm, Distal calf - 0.25cm

SPJ is competent. SSV is competent.