



Reason Claudication
Outcome Occlusion


Right



Good



Good



Good



Good

Brachial

Common Femoral

High Thigh

Low Thigh

Popiteal

High Calf

Peroneal

Anterior Tibial


Posterior Tibial

Dorsalis Pedis

Toe Pressure

Post Exercise

Left




Good



Good



Good



Good

Notes

BILATERAL LOWER LIMB ARTERIAL DUPLEX

RIGHT:

CFA - patent with moderate disease, good biphasic waveforms, PSV 217cm/s.

PFA origin - patent with good biphasic waveforms, PSV 118cm/s.

SFA - patent with mild to moderate disease, good biphasic waveforms, PSV 83-120cm/s.

PopA - patent with good biphasic waveforms, PSV 81-71cm/s.

TPT is patent. 3 run-off origins noted.

Assessed by Sharifa Kiyegga

Printed on 16/12/2019 at 2:44 am

Checked by



3690

ATA and pero A - patent along length, good biphasic waveforms at the ankle, PSV 87-89cm/s.
PTA - proximal to mid vessel appears patent. Distal vessel appears occluded to ankle.

LEFT:

CFA - patent with mild disease, good biphasic waveforms, PSV 118cm/s.

PFA origin - patent with good biphasic waveforms, PSV 87cm/s.

SFA - patent with mild to moderate disease, good biphasic waveforms, PSV 75-76cm/s.

PopA - patent with good biphasic waveforms, PSV 93cm/s.

TPT is patent. 3 run-off origins noted.

ATA and pero A - patent along length, good biphasic waveforms at the ankle, PSV 82-44cm/s.

PTA - appears patent at the origin, then vessel appears to occlude to the ankle.

Resting ABPI attempted, however PSV >200mmHG.

