



Reason Varicose vein
Outcome Competent

	Right		Left	
Deep Veins	Patency	Competency	Patency	Competency
Common Iliac Vein	Patent		Patent	
External Iliac Vein	Patent	Competent	Patent	Competent
Internal Iliac Vein				
Common Femoral Vein	Widely Patent	Competent	Widely Patent	Competent
Profunda Vein	Widely Patent	Competent	Widely Patent	Competent
Superficial Femoral Vein	Widely Patent	Competent	Widely Patent	Competent
Popliteal Vein	Widely Patent	Competent	Widely Patent	Competent
Posterior Tibial Vein	Widely Patent	Competent	Widely Patent	Competent
Anterior Tibial Vein	Widely Patent	Competent	Widely Patent	Competent
Peroneal Vein	Widely Patent	Competent	Widely Patent	Competent
Soleal Vein	Patent	Competent	Patent	Competent
Gastrocnemius	Patent	Competent	Patent	Competent
Superficial Veins				
Saphenofemoral Junction	Patent	Competent	Patent	Competent
L Saphenous Vein Above	Patent	Competent	Patent	Competent
L Saphenous Vein Below	Patent	Competent	Patent	Competent
Vein of Giacomini	Patent	Competent	Patent	Competent
Saphenopopiteal Junction	Patent	Competent	Not Identified	
S Saphenous Vein	Patent	Competent	Patent	
Evidence of D.V.T.				
Above the knee	No		No	
Popliteal	No		No	
Below the knee	No		No	

Notes

BILATERAL LOWER LIMB VENOUS DUPLEX ASSESSMENT

The IVC, CIV and EIV appear patent with good/reasonable wall-to-wall colour-filling, bilaterally (the EIV's appear competent but the CIV's could not be viewed with the patient standing and were assessed supine). Flow in the common femoral vein is phasic with respiration and responds normally to a Valsalva manoeuvre, suggesting proximal vein patency. All visualised deep veins appear widely patent and competent with no evidence of previous DVT, bilaterally.

RIGHT:

Sapheno-femoral junction (SFJ) is competent. Long Saphenous vein (LSV) is competent along its length.

Short Saphenous vein (SSV) is patent and competent along its length and is continuous with a competent

Assessed by Sharifa Kiyegga

Printed on 24/12/2019 at 3:22 am

Checked by



vein of Giacomini.

Sapheno-popliteal junction (SPJ) is patent and competent.

Lateral and posterior superficial thigh veins noted, however unable to locate source.

LEFT:

Sapheno-femoral junction (SFJ) is competent. Long Saphenous vein (LSV) is competent along its length.

Short Saphenous vein (SSV) is competent along its length and is continuous with a competent vein of Giacomini.

Sapheno-popliteal junction (SPJ) was not identified.

Small calibre anterior thigh veins noted, however unable to locate source.