



Patient

NHS No

D.O.B.

Patient Ref

Reason

Varicose vein

Outcome

DVT negative, Competent

	Right		Left	
Deep Veins	Patency	Competency	Patency	Competency
Common Iliac Vein				
External Iliac Vein				
Internal Iliac Vein				
Common Femoral Vein	Widely Patent	Competent	Widely Patent	Competent
Profunda Vein	Widely Patent	Competent	Widely Patent	Competent
Superficial Femoral Vein	Widely Patent	Competent	Widely Patent	Competent
Popliteal Vein	Widely Patent	Competent	Widely Patent	Competent
Posterior Tibial Vein	Widely Patent	Competent	Widely Patent	Competent
Anterior Tibial Vein	Widely Patent	Competent	Widely Patent	Competent
Peroneal Vein	Widely Patent	Competent	Widely Patent	Competent
Soleal Vein				
Gastrocnemius	Widely Patent	Competent	Widely Patent	Competent
Superficial Veins				
Saphenofemoral Junction	Widely Patent	Competent	Widely Patent	Competent
L Saphenous Vein Above	Widely Patent	Competent	Widely Patent	Competent
L Saphenous Vein Below	Widely Patent	Competent	Widely Patent	Competent
Vein of Giacomini	Not Identified		Widely Patent	Competent
Saphenopopiteal Junction	Widely Patent	Competent	Not Identified	
S Saphenous Vein	Widely Patent	Competent	Widely Patent	Competent
Evidence of D.V.T.				
Above the knee	No		No	
Popliteal	No		No	
Below the knee	No		No	

Notes**BILATERAL LOWER LIMB VENOUS DUPLEX ASSESSMENT**

Iliac veins not viewed, bilaterally. Flow in the right and left common femoral veins is phasic with respiration and a normal response on Valsalva manoeuvre, suggesting proximal vein patency. All visualised deep veins appear widely patent and competent with no evidence of previous, bilateral DVT.

All truncal, superficial veins appear to be widely patent and competent.

Assessed by

Lukasz Koprowski

Checked by