



Patient

NHS No

D.O.B.

Patient Ref

**Reason**

Varicose vein

**Outcome**

Incompetence

	<b>Right</b>		<b>Left</b>	
<b>Deep Veins</b>	<b>Patency</b>	<b>Competency</b>	<b>Patency</b>	<b>Competency</b>
Common Iliac Vein				
External Iliac Vein				
Internal Iliac Vein				
Common Femoral Vein			<input type="text" value="Widely Patent"/>	<input type="text" value="Competent"/>
Profunda Vein				
Superficial Femoral Vein				
Popliteal Vein			<input type="text" value="Widely Patent"/>	<input type="text" value="Competent"/>
Posterior Tibial Vein				
Anterior Tibial Vein				
Peroneal Vein				
Soleal Vein				
Gastrocnemius				
<b>Superficial Veins</b>				
Saphenofemoral Junction			<input type="text" value="Widely Patent"/>	<input type="text" value="Incompetent"/>
L Saphenous Vein Above			<input type="text" value="Widely Patent"/>	<input type="text" value="Isolated Incompetence"/>
L Saphenous Vein Below			<input type="text" value="Widely Patent"/>	<input type="text" value="Isolated Incompetence"/>
Vein of Giacomini			<input type="text" value="Not Identified"/>	
Saphenopopiteal Junction			<input type="text" value="Widely Patent"/>	<input type="text" value="Competent"/>
S Saphenous Vein			<input type="text" value="Widely Patent"/>	<input type="text" value="Competent"/>
<b>Evidence of D.V.T.</b>				
Above the knee			<input type="text" value="No"/>	
Popliteal			<input type="text" value="No"/>	
Below the knee				

**Notes****LEFT LOWER LIMB VENOUS DUPLEX ASSESSMENT**

Iliac veins not viewed. Flow in the common femoral vein is phasic with respiration and a normal response on Valsalva manoeuvre, suggesting proximal vein patency. Common femoral and popliteal veins are widely patent and competent with no evidence of DVT.

Sapheno-femoral junction (SFJ) is incompetent. Long saphenous vein (LSV) is incompetent and linear down to mid thigh. Incompetent branches leave the LSV in the mid thigh and forms the visible varicosities of the medial and posterior thigh. The LSV is competent distal to the afore mentioned branches, with the exception of the short section at the knee level (see diagram).

An incompetent perforator was noted in the mid-lateral thigh, forming the visible varicosities of the lateral thigh. The tributaries run inferiorly and confluence the visible varicosities originating from the LSV in the

Assessed by

Lukasz Koprowski

Checked by



Patient

NHS No


D.O.B.

Patient Ref

posterior, proximal calf.

Transverse (AP) dimensions of LSV: proximal thigh 0.67cm, mid thigh 0.62cm, knee level 0.37cm.

Sapheno-popliteal junction (SPJ) and short saphenous vein (SSV) are competent.



Assessed by

Lukasz Koprowski

Checked by