



Patient

NHS No

D.O.B.

Patient Ref

Reason

Varicose vein

Outcome

DVT negative, Incompetence

	Right		Left	
Deep Veins	Patency	Competency	Patency	Competency
Common Iliac Vein				
External Iliac Vein				
Internal Iliac Vein				
Common Femoral Vein	Widely Patent	Competent		
Profunda Vein	Widely Patent			
Superficial Femoral Vein	Widely Patent			
Popliteal Vein	Widely Patent	Competent		
Posterior Tibial Vein	Widely Patent			
Anterior Tibial Vein	Widely Patent			
Peroneal Vein	Widely Patent			
Soleal Vein				
Gastrocnemius	Patent			
<b>Superficial Veins</b>				
Saphenofemoral Junction	Widely Patent	Incompetent		
L Saphenous Vein Above	Widely Patent	Isolated Incompetence		
L Saphenous Vein Below	Widely Patent	Competent		
Vein of Giacomini	Widely Patent	Competent		
Saphenopopliteal Junction	Not Identified			
S Saphenous Vein	Widely Patent	Competent		
<b>Evidence of D.V.T.</b>				
Above the knee	No			
Popliteal	No			
Below the knee	No			

**Notes****RIGHT LOWER LIMB VENOUS DUPLEX ASSESSMENT**

Iliac veins not viewed. Flow in the common femoral vein is phasic with respiration and a normal response on Valsalva manoeuvre, suggesting proximal vein patency. All the deep veins, including the common femoral and popliteal veins are widely patent / patent and competent with no evidence of DVT, at this time.

Sapheno-femoral junction (SFJ) is incompetent. Long saphenous vein (LSV) is incompetent and linear in the thigh. Incompetent branch leaves the LSV in the distal thigh and forms the visible varicosities of the thigh and calf. Distal to this branch the LSV is competent.

Transverse (AP) dimensions of LSV: proximal thigh 0.75cm, mid-distal thigh 0.78cm.

Sapheno-popliteal junction (SPJ) was not identified. Short saphenous vein (SSV) is competent and is

Assessed by

Lukasz Koprowski

Checked by




Patient

NHS No

D.O.B.

Patient Ref

continuous with a competent vein of Giacomini.



Assessed by

Lukasz Koprowski

Checked by