



Patient	NHS No
D.O.B.	Patient Ref
Reason	Varicose vein
Outcome	?infection, DVT negative, Lymph nodes, Incompetence

	Right		Left	
Deep Veins	Patency	Competency	Patency	Competency
Common Iliac Vein				
External Iliac Vein				
Internal Iliac Vein				
Common Femoral Vein	Widely Patent	Competent		
Profunda Vein	Widely Patent	Competent		
Superficial Femoral Vein	Widely Patent	Competent		
Popliteal Vein	Widely Patent	Competent		
Posterior Tibial Vein	Widely Patent	Competent		
Anterior Tibial Vein	Widely Patent	Competent		
Peroneal Vein	Widely Patent	Competent		
Soleal Vein				
Gastrocnemius	Widely Patent	Competent		
Superficial Veins				
Saphenofemoral Junction	Widely Patent	Incompetent		
L Saphenous Vein Above	Widely Patent	Incompetent		
L Saphenous Vein Below	Widely Patent	Incompetent		
Vein of Giacomini	Widely Patent	Competent		
Saphenopopiteal Junction	Not Identified			
S Saphenous Vein	Widely Patent	Competent		
Evidence of D.V.T.				
Above the knee	No			
Popliteal	No			
Below the knee	No			

Notes

RIGHT LOWER LIMB VENOUS DUPLEX ASSESSMENT

Iliac veins not viewed. Flow in the common femoral vein is phasic with respiration and a normal response on Valsalva manoeuvre, suggesting proximal vein patency. All visualised deep veins appear widely patent and competent with no evidence of previous DVT.

All measurements are proximal to the medial malleolus unless otherwise stated.

Sapheno-femoral junction (SFJ) is incompetent. Long saphenous vein (LSV) is incompetent throughout its length, being linear and intra-fascial down to proximal calf. An incompetent branch leaves the LSV in the prox-mid calf (~27cm), tracks inferiorly and forms the minimally visible varicosities of the distal calf / dorsum of the foot.

Assessed by	Lukasz Koprowski	Checked by	
-------------	------------------	------------	--



Patient

NHS No


D.O.B.

Patient Ref

Sapheno-popliteal junction (SPJ) was not identified. Short saphenous vein (SSV) is competent and is continuous with a competent vein of Giacomini.

Transverse (AP) dimensions of LSV: Proximal thigh 0.57cm, Mid thigh 0.5cm, Distal thigh 0.64cm, Proximal calf 0.58cm, Mid calf 0.45cm, Distal calf 0.47cm.

ADDITIONAL FINDING: An avascular, mixed echogenic mass noted in the right groin, measuring ~1.6 cm ML x ~0.8cm TS ?enlarged lymph node ?infection.



Assessed by

Lukasz Koprowski

Checked by