



Patient	NHS No
D.O.B.	Patient Ref

Reason	Varicose vein
Outcome	DVT negative, Incompetence

	Right		Left	
Deep Veins	Patency	Competency	Patency	Competency
Common Iliac Vein				
External Iliac Vein				
Internal Iliac Vein				
Common Femoral Vein			Widely Patent	Competent
Profunda Vein			Widely Patent	Competent
Superficial Femoral Vein			Widely Patent	Competent
Popliteal Vein			Widely Patent	Competent
Posterior Tibial Vein			Widely Patent	Competent
Anterior Tibial Vein			Widely Patent	Competent
Peroneal Vein			Widely Patent	Competent
Soleal Vein				
Gastrocnemius			Widely Patent	Competent
Superficial Veins				
Saphenofemoral Junction			Widely Patent	Incompetent
L Saphenous Vein Above			Widely Patent	Incompetent
L Saphenous Vein Below			Widely Patent	Isolated Incompetence
Vein of Giacomini			Not Identified	
Saphenopopliteal Junction			Widely Patent	Competent
S Saphenous Vein			Widely Patent	Competent
Evidence of D.V.T.				
Above the knee			No	
Popliteal			No	
Below the knee			No	

## Notes

### LEFT LOWER LIMB VENOUS DUPLEX ASSESSMENT

Iliac veins not viewed. Flow in the common femoral vein is phasic with respiration and a normal response on Valsalva manoeuvre, suggesting proximal vein patency. All visualised deep veins appear widely patent and competent with no evidence of previous DVT.

All measurements are proximal to the medial malleolus unless otherwise stated.

Sapheno-femoral junction (SFJ) is incompetent. Long saphenous vein (LSV) is incompetent down to proximal calf and is linear in the thigh. Three notable, incompetent branches identified in the proximal (~26cm), mid (~22cm) and distal calf (~12cm), all supplying the visible varicosities of the medial calf. The LSV is competent between the branches at ~26cm and ~22cm, is incompetent between branches at

Assessed by                      Lukasz Koprowski

Checked by                      \_\_\_\_\_



Patient


NHS No

D.O.B.

Patient Ref

~22cm and ~12cm, and is competent distal to the branch at ~12cm.  
Sapheno-popliteal junction (SPJ) and short saphenous vein (SSV) are competent.

Transverse (AP) dimensions of LSV: Proximal thigh 0.64cm, Mid thigh 0.56cm, Distal thigh 0.59cm,  
Proximal calf 0.57cm, Mid-distal calf 0.49cm.



Assessed by

Lukasz Koprowski

Checked by