



Patient

NHS No

D.O.B.

Patient Ref

Reason

Varicose vein

Outcome

DVT negative, Superficial thrombophlebitis, Incompetence

	Right		Left	
Deep Veins	Patency	Competency	Patency	Competency
Common Iliac Vein				
External Iliac Vein				
Internal Iliac Vein				
Common Femoral Vein	<input type="text" value="Widely Patent"/>	<input type="text" value="Competent"/>		
Profunda Vein	<input type="text" value="Widely Patent"/>	<input type="text" value="Competent"/>		
Superficial Femoral Vein	<input type="text" value="Widely Patent"/>	<input type="text" value="Competent"/>		
Popliteal Vein	<input type="text" value="Widely Patent"/>	<input type="text" value="Competent"/>		
Posterior Tibial Vein	<input type="text" value="Widely Patent"/>	<input type="text" value="Competent"/>		
Anterior Tibial Vein	<input type="text" value="Widely Patent"/>	<input type="text" value="Competent"/>		
Peroneal Vein	<input type="text" value="Widely Patent"/>	<input type="text" value="Competent"/>		
Soleal Vein				
Gastrocnemius	<input type="text" value="Widely Patent"/>	<input type="text" value="Competent"/>		
Superficial Veins				
Saphenofemoral Junction	<input type="text" value="Widely Patent"/>	<input type="text" value="Incompetent"/>		
L Saphenous Vein Above	<input type="text" value="Areas of Thrombus"/>	<input type="text" value="Fresh & mixed thrombus"/>		
L Saphenous Vein Below	<input type="text" value="Areas of Thrombus"/>	<input type="text" value="Fresh & mixed thrombus"/>		
Vein of Giacomini	<input type="text" value="Widely Patent"/>	<input type="text" value="Competent"/>		
Saphenopopiteal Junction	<input type="text" value="Not Identified"/>			
S Saphenous Vein	<input type="text" value="Widely Patent"/>	<input type="text" value="Competent"/>		
Evidence of D.V.T.				
Above the knee	<input type="text" value="No"/>			
Popliteal	<input type="text" value="No"/>			
Below the knee	<input type="text" value="No"/>			

Notes

RIGHT LOWER LIMB VENOUS DUPLEX ASSESSMENT

Iliac veins not viewed. Flow in the common femoral vein is phasic with respiration and a normal response on Valsalva manoeuvre, suggesting proximal vein patency. All visualised deep veins appear widely patent and competent with no evidence of previous DVT.

All measurements are proximal to the medial malleolus unless otherwise stated.

Sapheno-femoral junction (SFJ) is incompetent. Extensive sections of long saphenous vein (LSV) down to mid calf contain irregular, incompetent flow with non-occlusive superficial thrombophlebitis.

Multiple, incompetent branches leave the LSV in the prox-mid calf, forming the visible varicosities of the postero-medial calf. Some of the branches contain superficial thrombophlebitis. Distal to a large branch at

Assessed by

Lukasz Koprowski

Checked by



Patient

NHS No


D.O.B.

Patient Ref

mid calf (~16cm), the LSV becomes small and appears to be competent.

Sapheno-popliteal junction (SPJ) was not identified. Short saphenous vein (SSV) is competent and is continuous with a competent vein of Giacomini.

An incompetent, superficial perforator noted in the posterior mid calf (~21cm).



Assessed by

Lukasz Koprowski

Checked by