

Ref. Locn. : **GENERAL SURGERY CLINIC BROOMFIELD**  
Referrer : **RAMIREZ MJUR**

DoB :  
Hosp. No. :  
CRIS No. :  
NHS No. :

**VERIFIED** Verified By: Ikhlq Ahmad Rana 01-Jun-2022 11:10 First Typed By: RQ8VIR 01-Jun-2022 1110.

Recurrent right Leg Varicose veins

Right:

Incompetent CFV, FV and Popliteal Vein drain to recanalised incompetent short saphenous vein from incompetent SPJ, diameter 7.6mm to 8.4mm.

There are scattered thrombi in recanalised Short saphenous vein.

Upper 10cm segment of incompetent SSV, communicate with obvious varicose veins on posterior calf and one main varicose vein, which swings anteromedially and then descends to foot.

SPJ is approximately 4cm above skin crease.

SSV is competent at mid and lower calf.

Long Saphenous Vein is competent.

crural veins are competent.

Scan performed and reported by: Ikhlq Rana

Event Number : E-7379030

Event Date/Time : **01-Jun-2022, 1029**

Ref. Source : RAMIREZ MJUR, Broomfield Hospital, Court Road, Chelmsford, Essex, CM1 7ET

Examinations : **US Doppler lower limb veins Rt**