

Ref. Locn. : **GENERAL SURGERY CLINIC BROOMFIELD**Referrer : **RAMIREZ MJUR**

DoB :

Hosp. No. :

CRIS No. :

NHS No. :

VERIFIED Verified By: Ikhlq Ahmad Rana 09-Mar-2022 0941 First Typed By: RQ8VIR 09-Mar-2022 0941.

Recurrent varicose veins.

Right;

GSV is thrombosed in the thigh.

A tortuous incompetent vein diameter 7.9mm to 9.0mm, arises from incompetent perforator 15cm below groin, also joined by small descending veins from groin, goes downward to medial knee, drain in to an incompetent at popliteal crease and gives rise to obvious varicose veins on anterior and posterior calf.

SSV and deep veins are patent and competent.

Left;

Incompetent recurrent veins along side thrombosed GSV arise from incompetent SFJ, join to form main tortuous vein, diameter 9mm, which gives rise to obvious varicose veins on medial anterior and lateral thigh+Anterior, lateral and posterior calf.

SSV and deep veins are patent and competent.

Scan performed and reported by: Ikhlq Rana

Event Number : E-7318750

Event Date/Time : **09-Mar-2022, 0849**

Ref. Source : RAMIREZ MJUR, Broomfield Hospital, Court Road, Chelmsford, Essex, CM1 7ET

Examinations : **US Doppler lower limb veins Both**