

**CL1.3****3. Extra-cranial carotid/ brachio-cephalic/ subclavian/ vertebral assessment**

Probe types – 12 - 3MHz

Measurements – Velocities in centimetres per second (cm/s), diameter (transverse; anterior-posterior, medial-lateral) in centimetres (cm) (if dilated/pre-op), length of disease (longitudinal) in cm.

Patient positioning and scanning approach – patients can be scanned supine or in a sitting position. A supine approach with the vascular scientist sat behind the patient's head allows easy access to the neck and reduces the risk of RSI (repetitive strain injury) as the operator can rest their arm on the pillow or on the head of couch. The patient extends the neck and turns the head in the opposite direction to the side being assessed. Both sides of the neck are always assessed¹.

The carotid arteries can be viewed from a lateral or antero-lateral approach using the sternocleidomastoid muscle as an acoustic window².

B-mode assessment

Intimal B-mode assessment is performed to achieve an accurate picture of the anatomy and identify the location of the carotid bifurcation as well as the presence of any plaque morphology^{2, 3}.

Using B-mode, the common carotid artery (CCA) should be imaged in cross-section (transverse plane) and traced proximally to the clavicle until the subclavian artery is visualised. The distal brachio-cephalic artery may be visualised on the right side of the neck. On the left side, the origin of the CCA and subclavian arteries will not be visualised due to depth. The CCA should then be scanned along its length to the level of the bifurcation where the internal carotid artery (ICA) and external carotid artery (ECA) are visualised from their origins as far distal as possible.

The same method should then be repeated in longitudinal plane².

Colourflow assessment

Using the Colourflow modality, the CCA is scanned longitudinally where it is traced from the proximal section at the level of the clavicle to the distal section where the bifurcation, ICA and ECA are visualised as far distal as possible.

Colour should be used to identify ECA branches, filling defects, occlusion and velocity changes/ turbulence, although diagnosis should not be made using colour Doppler alone^{2,3}.

Grading plaque morphology – greyscale echogenicity

Switching to the greyscale imaging mode, a note can be made of the site, type and extent of plaque morphology.

The subclavian is visualised along its length in longitudinal section. The CCA, ICA and ECA are then viewed in cross-section and longitudinally. As soft plaque has the same echogenicity as blood, colourflow is the best modality for identification.



Soft plaque – associated with higher lipid content or thrombus. May have an anechoic or echolucent appearance similar to that of blood/fluid^{2,3}

Mixed plaque – variable/ heterogenous appearance of mixed or random echoes with some echogenic and some echolucent areas^{2,3}.

Dense plaque – homogenous appearance of bright white echoes⁴.

Calcified plaque – acoustic shadowing cast from the hardened plaque^{2,3}.

Irregular – broken or irregular luminal surface but not generally an indication of ulceration¹⁶.

Ulcerative – an area of mixed plaque forming a ‘crater’ of at least 2mm depth. May be seen in cross-section as a ‘hook’ of mixed plaque surrounding soft plaque, or with blood visibly swirling within the crater^{2,3}.

Doppler assessment

In the absence of significant disease, peak systolic velocity (PSV) measurements are taken from the CCA (1-2cm proximal to bifurcation)^{1, 2}, ICA and ECA. If the peak velocities are raised above 1.3m/s then the end-diastolic velocity (EDV) is also measured.

If significant plaques have been identified using B-mode and colour flow Doppler then further spectral Doppler samples are taken to investigate velocity increases and analyse the degree of stenosis in particular vessel. Stenosis in the ICA is graded using the criteria explained below. Atypical waveform profiles should also be noted^{2, 3}.

In cross-section, the CCA is traced proximal towards the clavicle and the transducer is angled beneath the clavicle until the subclavian artery is viewed in longitudinal section. The subclavian is traced as far proximal and distal as possible making note of areas of turbulence or narrowing. The PSV is measured using Doppler ultrasound. A second Doppler reading is taken as far distal as possible and the waveform characteristics are recorded (e.g. triphasic, biphasic, monophasic, turbulent, damped etc.).

Velocities in kinked arteries are less reliable as vessel tortuosity can raise velocities¹⁷. Care must be taken to ensure that the angle is correct to blood flow rather than the vessel³. In reporting, it will be stated ‘peak velocities indicate x% - y% stenosis but no plaque morphology noted.

Grading degree of carotid stenosis

Normal Velocities:

ICA:

- average (avg) PSV = 54 – 88cm/s (distal to bulb)⁴
- avg PSV = 74cm/s, avg EDV = 29cm/s (distal to bulb)⁵
- velocity slightly elevated if patient hypertensive⁶
- maximum PSV noted in normal = 115cm/s⁷



ECA:

- avg PSV \approx 77cm/s (normally <115cm/s)⁴
- avg PSV = 84cm/s, avg EDV = 16cm/s⁵
- ECA velocities can be elevated by an ipsilateral ICA occlusion⁴

CCA:

- avg PSV = 60 – 100cm/s⁸
- avg PSV = 108 +/- 18 cm/s (mean +/- S.D.)⁹
- avg PSV = 78-108 cm/s⁷
- avg PSV = 99cm/s, avg EDV = 27cm/s⁵
- on average, PSV in L CCA exceeds PSV in R CCA by 5cm/s⁹
- velocity slightly elevated if patient hypertensive⁶

Carotid Criteria

Diameter Stenosis	Morphology	ICA PSV (cm/s)	ICA EDV (cm/s)	PSVica/PSVcca	St Mary's ratio
<25%	Normal	<125	<40		
<30%	Intimal Thickening	<125	<40		
<30%	Plaque	<125	<40		
<40%	Plaque	<125	<40		
<50%	Plaque	<125	<40	<2	<8
50-59%	Plaque	>125	<40	<3.2	8.0-10
60-69%	Plaque	>125	40-110	3.2-4.0	11-13
70-79%	Plaque	>230	110-140	>4.0	14-21
80-89%	Plaque	>230	>140	>4.0	22-29
90-95%	Plaque	>400	>140	>5.0	>30
96-99%	Plaque	Trickle flow			Variable
100%	Plaque	Absence of flow			N/A

Sidhu and Allan. Ultrasound Assessment of Internal Carotid Artery Stenosis. Clinical Radiology, (1997) 52, 654-658. (Developed using data from Moneta et al. 1993, 1995). CP Oates et al. Joint recommendations for Reporting Carotid Ultrasound Investigations in the UK. EurJ Vasc Endovasc Surg (2008) 20, 1-11.

Criteria are only reliable for internal carotid artery stenosis³.

ICA peak systolic velocities are less reliable in the presence of CCA disease and ratios should be used. The use of the ICA: CCA PSV ratio normalises ICA PSV measurements^{2 3}.

Elevated velocities can be produced in the CCA, ICA¹⁹ and ECA in the presence of contralateral CCA or ICA stenosis or occlusion.

A significant proximal (CCA origin or brachio-cephalic) ipsilateral stenosis can reduce velocities in the CCA, ICA and ECA.



Aortic stenosis can reduce the velocities in the CCA only.

Peak systolic velocities from large carotid bulbs may be unreliable, estimate degree of stenosis using grey scale and diameter/area reduction measurement.

Doppler Waveforms:

1. CCA waveform has a low-resistance pattern (most of the CCA flow goes to the brain). Note that a small amount of post systolic flow reversal (giving rise to a triphasic waveform) is normal; reversal of flow evident for more than 50% of the duration of diastole should be regarded as abnormal (see point 5 below)¹⁰.
2. Normal ICA waveform has low-resistance pattern (all of the ICA flow goes to brain)¹⁸.
3. Normal ECA waveform has a high-resistance pattern (vessel supplies a high resistance vascular bed). Note the prominent diastolic notch, which represents closure of the aortic valve and the onset of diastole¹⁰.
4. Severe proximal stenosis (innominate artery, CCA origin, aortic valve) produces a damped waveform (“tardus-parvus”, where tardus infers the pulse is slow to rise and fall and parvus infers a small pulse.)^{4,8}. **Essentially, the acceleration time to systole is increased, hence the slope of the systolic upstroke is reduced, and there is blunting and smoothing of the sharp peak representing a reduction in waveform pulsatility⁹.** This effect is usually most prominent in the CCA, but is also sometimes seen in the ICA & ECA. Note that in the case of aortic valve disease or diminished cardiac output, damping is symmetrical (seen in both CCAs)⁴.
5. Severe aortic incompetence with or without the presence of significant aortic stenosis often produces either a bisferious (two systolic peaks, well separated from the diastolic notch, with the second peak being the same height as or higher than the first) waveform¹⁰, or persistent reverse diastolic flow in the CCA, or both. Note that these effects are not usually seen in the ICA, but are evident in both the CCA & ECA.
6. Significant stenosis or occlusion of the distal CCA or the ICA causes a high-resistance ipsilateral CCA waveform; reverse flow is evident and often there is complete loss of end diastolic flow. Note that significant ECA disease does not usually impact on the CCA waveform due to its relatively low flow volume⁴.

External Carotid Artery Assessment

From searching the literature (pubmed, medline, science direct, quest) there is no evidence of a radiologically validated method for grading ECA disease using a velocity criteria.

There is normally little requirement for the grading of ECA disease due to its highly branched vascular network and non-cerebral involvement^{13,15}. In cases where a patient experiences cerebral or ocular symptoms in the presence of ipsilateral ICA occlusion it may be useful to grade and characterise ECA disease as a possible cause of emboli and transient ischaemic attack (TIA). There is much published evidence extolling the



benefit of surgical or radiological intervention for the treatment of ECA disease where there is ipsilateral ICA occlusion and a thorough examination of disease is important in these cases^{13,14,15}.

At present staff use a visual estimation and/or use of electronic callipers to measure degree and extent of stenotic disease.

In the presence of an ICA occlusion, electronic callipers should be used in the transverse and longitudinal planes to measure degree of ECA stenosis. Length of stenosis, plaque characterisation and degree of turbulence should also be recorded in the report.

Vertebral artery assessment

The vertebral artery (VA) can be viewed if the transducer is angled posterior. The flow direction should be the same as the carotid flow direction and is checked using the colourflow, **but more** importantly the Doppler sample volume. Vertebral flow is graded as orthograde, oscillatory (i.e. reversed in either systole or diastole alone) or retrograde^{2,3}. If no colourflow is identified within the vessel lumen – use spectral or power Doppler to investigate as it is more sensitive than colourflow⁴.

NORMAL VELOCITIES:

- avg PSV = 20-40 cm/s^{2,3}
- PSV<10cm/s should be regarded as potentially abnormal⁴
- Higher velocities may be normal in the dominant VA of an asymmetric pair.^{2,3}
- Higher velocities may be normal with contralateral VA occlusion.^{2,3}
-

DOPPLER WAVEFORMS:

1. Normal VA waveform has a low-resistance pattern (supplying the brain), with cephalad flow throughout the cycle^{2,4}.
2. If the VA has a high-resistance, antegrade (cephalic) flow pattern, there is probably a significant obstruction distal to the site of examination. (The second most common site of VA atheroma is intracranially, just beyond the C1 arch)³.
3. Severe proximal stenosis produces a damped waveform; note that the most common site of VA atheroma is the VA origin, although this can be difficult to image as it originates from the posterior aspect of the subclavian artery³.
4. Subclavian artery origin stenosis can have varying effects on the VA waveform shape and the direction of flow, dependent on the degree of stenosis and the presence of other collateral pathways.



Pre-operative carotid assessment.

Staff must follow additional criteria when performing a pre-operative scan for carotid endarterectomy.

- 1.Length of disease from the bifurcation, into the ICA, must be documented.
- 2.Bifurcation needs to be marked on the skin surface – the image of the bifurcation is obtained then the probe is moved until the bifurcation is just off the leading edge of the probe, marks are made on the skin surface in transverse and longitudinal section. Where these lines transverse is the position of the bifurcation and an arrow should be drawn to mark the tip.
- 3.Take a picture of the disease and keep with our hardcopy.
- 4.Mark MCA signal – see TCD section

REFERENCES:

1. CP Oates et al. Joint Recommendations for Reporting Carotid Ultrasound Investigations in the United Kingdom. Eur J Vasc Endovasc Surg (2008) 20, 1-11.
2. Thrush and Hartshorn Vascular Ultrasound, How, Why and When. 3rd Ed 2010
3. Vascular Laboratory Practice Part II, IPEM 1st Ed 2001.
4. Zwiebel WJ. Introduction to Vascular Sonography, 4th Ed. 2000.
5. Halpern EJ, Merton DA, Forsberg F. Effect of distal resistance on Doppler US flow patterns. Radiology 1998; 206:761-766.
6. Ferrara LA, Mancini M, Iannuzzi R, Marotta T, Gaeta I, Pasanisi F, Postiglione A, Guida L. Carotid diameter and blood flow velocities in cerebral circulation in hypertensive patients. Stroke, 1995;26:418-421.
7. Meyer JJ, Khalil RM, Obuchowski NA, Baus LK. Common carotid artery: variability of Doppler US velocity measurements. Radiology 1997;204:339-341.
8. O'Boyle MK, Vibhakar NI, Chung J, Keen WD, Gosink BB. Duplex sonography of the carotid arteries in patients with isolated aortic stenosis: image findings and relation to severity of stenosis. AJR, 1996;166:197-202.
9. Holdsworth DW, Norley CJD, Frayne R, Steinman DA, Rutt BK. Characterisation of common carotid artery blood-flow waveforms in normal human subjects. Physiological Measurements, 1999;20:219-240.
10. Kallman CE, Gosnik BB, Gardner DJ. Carotid Duplex sonography: bisferious pulse contour in patients with aortic valvular disease. AJR, 1991; 157:403-407.
11. Malaterre HR, Kallee K, Giusiano B, Letallec L, Djiane P. Holodiastolic reversal flow in the common carotid: another indicator of the severity of aortic regurgitation. International Journal of Cardiovascular Imaging, 2001; 17(5): 333-337.
12. iu22 Ultrasound System. Getting Started handbook. Philips, Sept 2005.
13. Louvelos, GN et al. (2012) 'Endovascular repair of external carotid artery disease'. Journal of endovascular therapy.19 (4), 504-11
14. Eisenberg, J et al. (2005) 'Endovascular repair of symptomatic external carotid artery stenosis'. Journal of vascular surgery. 42(6), 1210–1212
15. Dwivedi AJ et al. (2011) 'Symptomatic external carotid artery stenosis' The American Surgeon. 77(11) pE238-239.



16. Prabhakaran, S. et al. (2006) 'Carotid Plaque Surface Irregularity Predicts Ischemic Stroke: The Northern Manhattan Study'. Stroke 37 (11), 2696-2701
17. Gaitini, D. and Soudack, M. (2005) 'Diagnosing Carotid Stenosis by Doppler Sonography'. JUM 24 (8), 1127-1136
18. Rohren, Eric M. et al. (2003) 'A Spectrum Of Doppler Waveforms In The Carotid And Vertebral Arteries'. American Journal of Roentgenology 181 (6), 1695-1704.
19. Henderson RD. et al. (2000) 'Effect of Contralateral Carotid Artery Stenosis on Carotid Ultrasound Velocity Measurements'. Stroke. 31, 2636-2640