



varicose vein post-op.

## Outcome

Superficial thrombophlebitis, Poor images, patient habitus, Incompetence

	Right		Left	
Deep Veins	Patency	Competency	Patency	Competency
Common Iliac Vein				
External Iliac Vein				
Internal Iliac Vein				
Common Femoral Vein			Patent	
Profunda Vein				
Superficial Femoral Vein				
Popliteal Vein				
Posterior Tibial Vein				
Anterior Tibial Vein				
Peroneal Vein				
Soleal Vein				
Gastrocnemius				
Superficial Veins				
Saphenofemoral Junction			Patent	
L Saphenous Vein Above			Occluded	Mixed Thrombus
L Saphenous Vein Below			Patent	Isolated Incompetence
Vein of Giacomini				
Saphenopopliteal Junction				
S Saphenous Vein				
Evidence of D.V.T.				
Above the knee			No	
Popliteal				
Below the knee				

## Notes

LEFT LOWER LIMB SUPERFICIAL VENOUS ASSESSMENT - POST LSV RFA

\*Challenging assessment due to patient body habitus and vessel depth\*

The left common femoral is widely patent and competent, with no evidence of DVT.  
The SFJ appears patent.

The LSV appears occluded and incompressible in the thigh, with no evidence of flow seen in vessel lumen.

The LSV is patent and competent in the very distal thigh and proximal calf. Medial calf and anterior shin varicosities still present. The LSV is competent to the mid calf before a communicating incompetent branch from the incompetent varicosities render it incompetent, which it remains to the ankle.

Assessed by Rae Larmour

Printed on 16/06/2022 at 1:25 pm

Checked by