



Reason	Pre-op CABG
Outcome	disease - mild

Right	Diameter (cm)	PSV (m/s)	EDV (m/s)	Stenosis
Common		0.35		< 30%
Plaque	Mixed			
Disease length from BIF				
Bifurcation				< 40%
Plaque	Dense Mixed			
Disease length from BIF				
Internal		0.56		< 30%
Plaque	Dense			
Disease length from BIF				
	Pk ICA/Pk CCA = 1.6			
External		0.50		< 30%
Plaque	Dense Mixed			
Disease length from BIF				
Vertebral	Open Orthograde			
Subclavian	No Turbulence	Good Signal	Triphasic	Widely Patent

Left	Diameter (cm)	PSV (m/s)	EDV (m/s)	Stenosis
Common		0.62		< 30%
Plaque	Mixed			
Disease length from BIF				
Bifurcation				< 30%
Plaque	Mixed			
Disease length from BIF				
Internal		0.48		< 30%
Plaque	Mixed			
Disease length from BIF				
	Pk ICA/Pk CCA = 0.8			
External		0.74		< 30%
Plaque	Mixed			
Disease length from BIF				
Vertebral	Open Orthograde			
Subclavian	No Turbulence	Good Signal	Triphasic	Widely Patent

Stenosis based on NASCET methods.

Disease within large diameter carotid bulb is measured using direct diameter methods as recommended in Oates et al (2009).

Notes

Mixed plaques identified in the right and left internal carotid artery, forming a less than 30% stenosis, bilaterally.

Assessed by Ranit Shail, MCVS

Printed on 24/07/2024 at 1:33 pm

Checked by _____

Please note, this is a technical report to be interpreted by a medical professional. If you are a patient reading the report and require further help, please discuss the report with the person who referred you for the examination.