

**Carotid Duplex****Sheikh Muhammad Saleem**Examined **24/12/2018 09:45**

Page 1 of 1

Reference

Accession **CR-18-0066831**Patient **Gillian Patterson**NHS No **498 893 8549**D.O.B. **17/04/1946**Patient Ref **5332862****Reason** TIA**Outcome** Intimal thickening

Right	Diameter (cm)	PSV (m/s)	EDV (m/s)	Stenosis
Common		0.78		< 30%
Plaque	Intimal Thickening			
Disease length from BIF				
Bifurcation				< 30%
Plaque	Intimal Thickening			
Disease length from BIF				
Internal		0.52		< 30%
Plaque	Intimal Thickening			
Disease length from BIF				
	Pk ICA/Pk CCA = 0.7			
External		0.81		< 30%
Plaque	Intimal Thickening			
Disease length from BIF				
Vertebral	Open Orthograde			
Subclavian	No Turbulence	Good signal	Triphasic	Widely Patent

Left	Diameter (cm)	PSV (m/s)	EDV (m/s)	Stenosis
Common		1.09		< 25%
Plaque	Normal			
Disease length from BIF				
Bifurcation				< 25%
Plaque	Normal			
Disease length from BIF				
Internal		0.87		< 25%
Plaque	Normal			
Disease length from BIF				
	Pk ICA/Pk CCA = 0.8			
External		0.73		< 25%
Plaque	Normal			
Disease length from BIF				
Vertebral	Open Orthograde			
Subclavian	No Turbulence	Good signal	Triphasic	Widely Patent

Stenosis based on NASCET velocity criteria.

Joint recommendations for reporting carotid ultrasound investigations in the United Kingdom'. Oates et al. Eur J Vasc Endovasc Surg. 2009 Mar;37(3):251-61

Notes**CAROTID DUPLEX**

Intimal thickening identified in the right internal carotid artery, forming a less than 30% reduction in luminal diameter.

The left internal carotid arteries appear widely patent with no evidence of any plaque morphology, intimal dissection or other abnormality identified.

Assessed by **Sharifa Kiyegga**

Printed on 25/01/2019 at 4:34 pm

Checked by _____