

**Varicose Vein****Gareth Harrison**Examined **01/11/2018 10:40**

Page 1 of 2

Reference

Accession **CR-18-0057312****Patient** **Lisa Oakes**
D.O.B. **22/10/1970****NHS No** **610 714 9996**
Patient Ref **5053840****Reason** Varicose vein
Outcome Incompetence

	Right		Left	
Deep Veins	Patency	Competency	Patency	Competency
Common Iliac Vein				
External Iliac Vein				
Internal Iliac Vein				
Common Femoral Vein	Widely Patent	Competent		
Profunda Vein	Widely Patent	Competent		
Superficial Femoral Vein	Widely Patent (scarring)	Competent		
Popliteal Vein	Widely Patent	Competent		
Posterior Tibial Vein	Widely Patent	Competent		
Anterior Tibial Vein	Widely Patent	Competent		
Peroneal Vein	Widely Patent	Competent		
Soleal Vein	Widely Patent	Competent		
Gastrocnemius	Widely Patent	Competent		

Superficial Veins

Saphenofemoral Junction	Not Assessed	
L Saphenous Vein Above	see notes	
L Saphenous Vein Below	Patent	Isolated Incompetence
Vein of Giacomini	Widely Patent	Competent
Saphenopopiteal Junction	Not Identified	
S Saphenous Vein	Patent	Competent

Evidence of D.V.T.

Above the knee	No
Popliteal	No
Below the knee	No

Notes**RIGHT LOWER LIMB VENOUS DUPLEX ASSESSMENT**

Iliac veins not viewed. Flow in the common femoral vein is phasic with respiration and responds normally to a Valsalva manoeuvre, suggesting proximal vein patency. All visualised deep veins appear widely patent and competent. Isolated venous scarring identified in the distal SFV. No evidence of previous DVT in other deep veins.

All measurements are proximal to the medial malleolus unless otherwise stated.

Sapheno-femoral junction (SFJ) is not identified. A junction with an incompetent anterior thigh vein (ATV) at the CFV was identified at the 12 O'clock. The ATV forms small incompetent and tortuous veins in the anterior thigh ?neovascularisation. Long Saphenous vein (LSV) was not identified in the proximal to mid

Assessed by **Sharifa Kiyegga**

Printed on 24/12/2018 at 5:58 pm

Checked by _____



Patient **Lisa Oakes**
D.O.B. **22/10/1970**

NHS No **610 714 9996**
Patient Ref **5053840**

thigh. An incompetent LSV reforms in the very distal thigh at 48cm at which point an incompetent branch leaves the fascia. An incompetent branch leaves the fascia at knee level at 41.5cm forming varicosities. Distal to this the LSV is competent until an incompetent branch communicates with in the LSV in the mid calf at 22cm. LSV remains incompetent to the ankle (AP diameter - 0.32cm - 0.26cm).

Sapheno-popliteal junction (SPJ) was not identified.

Short Saphenous vein (SSV) is competent and is continuous with a competent vein of Giacomini.

