



Patient

Dale Clare

NHS No

612 865 7044

D.O.B.

07/06/1971

Patient Ref

3004274

Reason

Varicose vein

Outcome

Incompetence

	Right		Left	
Deep Veins	Patency	Competency	Patency	Competency
Common Iliac Vein				
External Iliac Vein				
Internal Iliac Vein				
Common Femoral Vein			Widely Patent	Competent
Profunda Vein			Widely Patent	Competent
Superficial Femoral Vein			Widely Patent	Competent
Popliteal Vein			Widely Patent	Competent
Posterior Tibial Vein			Widely Patent	Competent
Anterior Tibial Vein			Widely Patent	Competent
Peroneal Vein			Widely Patent	Competent
Soleal Vein			Widely Patent	Competent
Gastrocnemius			Widely Patent	Competent
Superficial Veins				
Saphenofemoral Junction			Patent	Incompetent
L Saphenous Vein Above			Patent	see notes
L Saphenous Vein Below			Patent	Isolated Incompetence
Vein of Giacomini			Patent	Competent
Saphenopopiteal Junction			Patent	Competent
S Saphenous Vein			Patent	Competent
Evidence of D.V.T.				
Above the knee			No	
Popliteal			No	
Below the knee			No	

Notes

LEFT LOWER LIMB VENOUS DUPLEX ASSESSMENT:

*Please noted diagram is of right leg but is actually left leg.

Iliac veins not viewed. Flow in the common femoral vein is phasic with respiration and responds normally to a Valsalva manoeuvre, suggesting proximal vein patency. All visualised deep veins appear widely patent and competent with no evidence of previous DVT.

All measurements are proximal to the medial malleolus unless otherwise stated.

Sapheno-femoral junction (SFJ) is incompetent. Long Saphenous vein (LSV) is incompetent in the proximal thigh (AP diameter: 0.75cm). The LSV appears to branch at 6cm distal to the groin with the more lateral

Assessed by Sharifa Kiyegga

Printed on 25/01/2019 at 4:48 pm

Checked by



Patient

Dale Clare

NHS No

612 865 7044

D.O.B.

07/06/1971

Patient Ref

3004274

branch remaining incompetent to the mid thigh (AP diameter: 0.48cm). The lateral branch leaves the fascia in the mid thigh at 61cm forming thigh varicosities.

Medial LSV: is competent to the very distal thigh. A small incompetent branch communicates with the LSV in the very distal thigh at 47cm. Distal to the the LSV is incompetent and remains incompetent until an incompetent branch leaves the fascia in the proximal calf at 33cm. Distal to this the LSV is competent until another incompetent branch communicates with it in the mid calf at 27cm. The LSV is incompetent distal to this and remains incompetent to the mid/distal calf. An incompetent branch leaves the fascia in the mid/distal calf at 18cm. Distal to this the LSV is competent to the ankle.

Short Saphenous vein (SSV) is competent and is continuous with a competent vein of Giacomini.

Saphno-popliteal junction is competent

