



Patient **Mary Scott**
D.O.B. **10/11/1927**

NHS No **440 380 5019**
Patient Ref **3073070**

Reason Varicose vein
Outcome Incompetence

	Right		Left	
	Patency	Competency	Patency	Competency
Deep Veins				
Common Iliac Vein				
External Iliac Vein				
Internal Iliac Vein				
Common Femoral Vein			Widely Patent	Competent
Profunda Vein			Widely Patent	Competent
Superficial Femoral Vein			Widely Patent	Isolated Incompetence
Popliteal Vein			Widely Patent	Isolated Incompetence
Posterior Tibial Vein			Patent	Competent
Anterior Tibial Vein			Patent	Competent
Peroneal Vein			Patent	Competent
Soleal Vein			Patent	Competent
Gastrocnemius			Patent	Competent
Superficial Veins				
Saphenofemoral Junction			Patent	Competent
L Saphenous Vein Above			Patent	Slight Incompetence
L Saphenous Vein Below			Patent	Slight Incompetence
Vein of Giacomini			Patent	
Saphenopopiteal Junction			Not Identified	
S Saphenous Vein			Patent	Competent
Evidence of D.V.T.				
Above the knee			No	
Popliteal			No	
Below the knee			No	

Notes**LEFT LOWER LIMB VENOUS DUPLEX ASSESSMENT**

Iliac veins not viewed. Flow in the common femoral vein is phasic with respiration and responds normally to a Valsalva manoeuvre, suggesting proximal vein patency. All visualised deep veins appear widely patent/patent with no evidence of previous DVT.. Isolated incompetence noted in the distal SFV and popliteal vein. All other deep vein appear competent.

All measurements are proximal to the medial malleolus unless otherwise stated.

Sapheno-femoral junction (SFJ) is competent. Slight incompetence noted in the Long Saphenous vein (LSV) in the proximal thigh. An incompetent branch leaves the fascia in the proximal/ mid thigh at 60cm forming thigh and calf varicosities. Distal to this the LSV is competent to the proximal calf. Isolated slight

Assessed by Sharifa Kiyegga

Printed on 01/02/2019 at 11:54 am

Checked by



Patient **Mary Scott**
D.O.B. **10/11/1927**

NHS No **440 380 5019**
Patient Ref **3073070**

incompetence noted in the calf LSV. Incompetent branches off the LSV noted in the proximal, mid and distal calf.

Transverse (AP) dimensions of LSV: Proximal thigh - 0.54-0.62cm.

Sapheno-popliteal junction (SPJ) was not identified.

Short Saphenous vein (SSV) is competent and is continuous with a competent vein of Giacomini.

