



Patient **Angela Maylor**
D.O.B. **19/07/1956**

NHS No **452 267 9823**
Patient Ref **3051715**

Reason Varicose vein
Outcome Incompetence

	Right		Left	
Deep Veins	Patency	Competency	Patency	Competency
Common Iliac Vein				
External Iliac Vein				
Internal Iliac Vein				
Common Femoral Vein	Widely Patent	Competent		
Profunda Vein	Widely Patent	Competent		
Superficial Femoral Vein	Widely Patent	Competent		
Popliteal Vein	Widely Patent	Competent		
Posterior Tibial Vein	Widely Patent	Competent		
Anterior Tibial Vein	Widely Patent	Competent		
Peroneal Vein	Widely Patent	Competent		
Soleal Vein	Patent	Competent		
Gastrocnemius	Patent	Competent		
Superficial Veins				
Saphenofemoral Junction	Patent	Slight Incompetence		
L Saphenous Vein Above	Patent	Isolated Incompetence		
L Saphenous Vein Below	Patent	Competent		
Vein of Giacomini	Not Identified			
Saphenopopiteal Junction	Patent	Competent		
S Saphenous Vein	Patent	Competent		
Evidence of D.V.T.				
Above the knee	No			
Popliteal	No			
Below the knee	No			

Notes

RIGHT LOWER LIMB VENOUS DUPLEX ASSESSMENT

Iliac veins not viewed. Flow in the common femoral vein is phasic with respiration and responds normally to a Valsalva manoeuvre, suggesting proximal vein patency. All visualised deep veins appear widely patent and competent with no evidence of previous DVT.

All measurements are proximal to the medial malleolus unless otherwise stated.

Sapheno-femoral junction (SFJ) is slightly incompetent. Long Saphenous vein (LSV) is incompetent in the proximal to mid thigh. An incompetent branch leaves the fascia in the mid thigh at 53cm, forming visible varicosities. Distal to the the LSV is competent to the ankle.

Assessed by Sharifa Kiyegga

Printed on 11/02/2019 at 3:21 pm

Checked by



Varicose Vein

Gareth Harrison

Examined **07/02/2019 10:40**

Page 2 of 2

Reference

Accession **CR-19-0007153**

Patient **Angela Maylor**


NHS No **452 267 9823**

D.O.B. **19/07/1956**

Patient Ref **3051715**

Transverse (AP) dimensions of LSV: Proximal thigh - 0.71cm, Mid- thigh - 0.34cm.

Sapheno-popliteal junction (SPJ) is patent and competent. SSV is competent along its length. Isolated venous scarring identified in the SSV.



Assessed by **Sharifa Kiyegga**

Printed on 11/02/2019 at 3:21 pm

Checked by _____